arXiv:1608.06885v1 [math-ph] 24 Aug 2016

QUANTUM DIMENSIONS AND FUSION PRODUCTS FOR IRREDUCIBLE V_Q^σ-MODULES WITH $\sigma^2 = 1$

JASON ELSINGER

Abstract. Every isometry σ of a positive-definite even lattice Q can be lifted to an automorphism of the lattice vertex algebra V_Q. An important problem in vertex algebra theory and conformal field theory is to classify the representations of the σ-invariant subalgebra V_Q^σ of V_Q, known as an orbifold. In the case when σ is an isometry of Q of order two, we have classified the irreducible modules of the orbifold vertex algebra V_Q^σ and identified them as submodules of twisted or untwisted V_Q-modules in [BE]. Here we calculate their quantum dimensions and fusion products.

The examples where Q is the orthogonal direct sum of two copies of the A_2 root lattice and σ is the 2-cycle permutation as well as where Q is the A_n root latice and σ is a Dynkin diagram automorphism are presented in detail.

1. Introduction

The notion of a vertex algebra has been a powerful tool for studying representations of infinite-dimensional Lie algebras. The theory of vertex algebras was introduced by Borcherds [B] and has since been developed in a number of works (see e.g. [FLM, K, FB, LL, KRR]).

Let σ be an automorphism of a vertex algebra V. An *orbifold* is the subalgebra of σ-invariants in V, denoted V^σ (see [DVVV, KT, DLM1]). It will follow from definitions below that every σ-twisted representation of V becomes untwisted when restricted to V^σ.

It has been a long-standing conjecture that all irreducible V^σ-modules are obtained by restriction from twisted or untwisted V-modules. This conjecture has recently been proved in a series of works by M. Miyamoto (see [M1, M2, M3, MC]) under suitable assumptions. In particular, it is shown in [MC] that if V is a simple regular vertex operator algebra of CFT type, then the fixed point subalgebra V^σ for a finite automorphism σ of V is also regular.

Date: August 25, 2016.

1991 *Mathematics Subject Classification.* Primary 17B69; Secondary 81R10.

Key words and phrases. Lattice vertex algebra; orbifold; twisted module; intertwining operator; quantum dimension; fusion product.

In this paper, we are concerned with the case when Q is a positive-definite even lattice and σ is an isometry of Q of order two. In [BE], we have classified and explicitly constructed all irreducible modules of the orbifold vertex algebra V_Q^σ, and have realized them as submodules of twisted or untwisted V_Q-modules. This paper is a continuation, where we give explicitly the quantum dimensions and fusion products for the irreducible V_Q^σ-modules.

Our general approach is to restrict from V_Q^σ to V_L^σ, where L is the sublattice of Q spanned by eigenvectors of σ. The subalgebra V_L^σ then factors as a tensor product $V_{L_+} \otimes V_{L_-}^\sigma$, where

$$L_\pm = \{\alpha \in Q \,|\, \sigma\alpha = \pm\alpha\},$$

and V_L^σ is rational by results of [DLM2, ABD, A2, DJL, FHL]. We then use the known irreducible V_L^σ-modules (see [DN, AD]) and their intertwining operators (see [A1, ADL]) to determine the quantum dimensions and fusion products for the irreducible V_Q^σ-modules given in [BE].

The paper is organized as follows. In Section 2, we briefly review lattice vertex algebras, their twisted modules, and the results for $\sigma = -1$ that we need. We also review the notions of quantum dimension and fusion product including results from [DJX] that we need. In Section 3, we review the irreducible orbifold modules given in [BE]. Our main results are presented in Sections 4 and 5. Examples where Q is the orthogonal direct sum of two copies of the A_2 root lattice and σ is the 2-cycle permutation as well as where Q is the A_n root latice and σ is a Dynkin diagram automorphism are presented in detail in Section 6.

2. Lattice Vertex algebras and their twisted modules

In this section, we briefly review lattice vertex algebras and their twisted modules, and recall the results in the case when $\sigma = -1$. These results will aid in the general description. General references on vertex algebras are e.g. [FLM, K, FB, LL, KRR], among many other works.

2.1. **Vertex algebras and twisted modules.**

A *vertex algebra* is a vector space V with a *vacuum vector* $\mathbf{1} \in V$ together with a linear map

$$Y(\cdot, z)\cdot \colon V \otimes V \to V[[z]][z^{-1}] \tag{2.1}$$

satisfying the axioms listed below. For $v \in V$, we have

$$Y(v, z) = \sum_{n\in\mathbb{Z}} v_{(n)} z^{-n-1}, \qquad v_{(n)} \in \operatorname{End} V. \tag{2.2}$$

The vacuum vector is the identity in the sense that

$$a_{(-1)}\mathbf{1} = \mathbf{1}_{(-1)}a = a\,, \qquad a_{(n)}\mathbf{1} = 0\,, \quad n \geq 0\,.$$

In particular,

$$\lim_{z\to 0} Y(a,z)\mathbf{1} = a.$$

For every $a \in V$, we call the image $Y(a,z)$ a *field*, meaning it can be viewed as a formal power series from $(\operatorname{End} V)[[z,z^{-1}]]$ involving only finitely many negative powers of z when applied to any vector.

The main axiom for a vertex algebra is the *Borcherds identity* (also called Jacobi identity in [FLM]) satisfied by the fields:

(2.3)
$$y^{-1}\delta\left(\frac{z_1-z_2}{y}\right)Y(u,z_1)Y(v,z_2) - y^{-1}\delta\left(\frac{z_2-z_1}{-y}\right)Y(v,z_2)Y(u,z_1)$$
$$= z_2^{-1}\delta\left(\frac{z_1-y}{z_2}\right)Y(Y(u,y)v,z_2),$$

where $u, v \in V$ and δ is the delta function $\delta(z) = \sum_{n\in\mathbb{Z}} z^n$. It is important to note that the commutator $[Y(u,z_1), Y(v,z_2)]$ follows from Borcherds identity be taking residue of both sides with respect to y.

A *V-module* for a vertex algebra V is a vector space M endowed with a linear map $Y^M(\cdot,z)\cdot\colon V\otimes M \to M[[z]][z^{-1}]$ (cf. (2.1)) such that the Borcherds identity (2.3) holds for all $u, v \in V$ which act as operators on M (see [FB, LL, KRR]). Let σ be an automorphism of V of finite order r. Then σ is diagonalizable. In the definition of a *σ-twisted V-module* M [FFR, D2], the image of Y^M is allowed to have nonintegral rational powers of z:

$$Y(u,z) = \sum_{n\in p+\mathbb{Z}} u_{(n)}\, z^{-n-1}\,, \qquad \text{if} \quad \sigma u = e^{-2\pi \mathrm{i} p}u\,,\ p \in \frac{1}{r}\mathbb{Z}\,,$$

where $u_{(n)} \in \operatorname{End} M$. The Borcherds identity in the twisted case is similar to (2.3), and it requires that u be an eigenvector of σ (see [FLM, FFR, D2]).

Recall from [FHL] that if V_1 and V_2 are vertex algebras, their tensor product is again a vertex algebra via

$$Y(v_1\otimes v_2, z) = Y(v_1,z)\otimes Y(v_2,z)\,, \qquad v_i \in V_i\,.$$

Furthermore, if M_i is a V_i-module, then the above formula defines the structure of a $(V_1\otimes V_2)$-module on $M_1\otimes M_2$ (see [FHL]). A similar statement is also true for twisted modules (see [BE, Lemma 2.2]).

2.2. **Lattice vertex algebras.** Consider an *integral lattice*, i.e., a free abelian group Q of finite rank together with a symmetric nondegenerate bilinear form $(\cdot|\cdot)\colon Q\times Q\to\mathbb{Z}$. We assume that Q is *even*, i.e., $|\alpha|^2=(\alpha|\alpha)\in 2\mathbb{Z}$ for all $\alpha\in Q$. The corresponding complex vector space $\mathfrak{h}=\mathbb{C}\otimes_{\mathbb{Z}}Q$ is considered as an abelian Lie algebra with a bilinear form extended from Q.

The *Heisenberg algebra* $\hat{\mathfrak{h}}=\mathfrak{h}[t,t^{-1}]\oplus\mathbb{C}K$ is the Lie algebra with brackets $[K,\mathfrak{h}]=0$, and

$$[a_m,b_n]=m\delta_{m,-n}(a|b)K\,,\qquad a_m=at^m\,. \tag{2.4}$$

Its induced irreducible highest-weight representation

$$M(1)=\mathrm{Ind}_{\mathfrak{h}[t]\oplus\mathbb{C}K}^{\hat{\mathfrak{h}}}\mathbb{C}\cong S(\mathfrak{h}[t^{-1}]t^{-1})$$

on which $K=1$ is known as the (bosonic) *Fock space*.

Following [FK, B], let $\varepsilon\colon Q\times Q\to\{\pm1\}$ be a bimultiplicative 2-cocycle such that

$$\varepsilon(\alpha,\alpha)=(-1)^{|\alpha|^2/2}\,,\qquad \alpha\in Q\,. \tag{2.5}$$

Then we have the associative algebra $\mathbb{C}_\varepsilon[Q]$ with basis $\{e^\alpha\}_{\alpha\in Q}$ and multiplication

$$e^\alpha e^\beta=\varepsilon(\alpha,\beta)e^{\alpha+\beta}\,. \tag{2.6}$$

Such a 2-cocycle ε is unique up to equivalence. In addition,

$$\varepsilon(\alpha,\beta)\varepsilon(\beta,\alpha)=(-1)^{(\alpha|\beta)}\,,\qquad \alpha,\beta\in Q\,. \tag{2.7}$$

The *lattice vertex algebra* associated to Q is defined as $V_Q=M(1)\otimes\mathbb{C}_\varepsilon[Q]$, where the vacuum vector is $\mathbf{1}=1\otimes e^0$. An action of the Heisenberg algebra on V_Q is given by

$$a_ne^\beta=\delta_{n,0}(a|\beta)e^\beta\,,\quad n\geq 0\,,\qquad a\in\mathfrak{h}\,.$$

The map Y on V_Q is uniquely determined by the generating fields:

$$Y(a_{-1}\mathbf{1},z)=\sum_{n\in\mathbb{Z}}a_n\,z^{-n-1}\,,\qquad a\in\mathfrak{h}\,, \tag{2.8}$$

$$Y(e^\alpha,z)=e^\alpha z^{\alpha_0}\exp\Big(\sum_{n<0}\alpha_n\frac{z^{-n}}{-n}\Big)\exp\Big(\sum_{n>0}\alpha_n\frac{z^{-n}}{-n}\Big)\,, \tag{2.9}$$

where $z^{\alpha_0}e^\beta=z^{(\alpha|\beta)}e^\beta$. Since the map $\mathfrak{h}\to M(1)$ given by $a\mapsto a_{-1}\mathbf{1}$ is injective, we can identify $a\in\mathfrak{h}$ with $a_{-1}\mathbf{1}\in M(1)$, i.e., $a_{(n)}=a_n$ for all $n\in\mathbb{Z}$.

2.3. **Twisted representations of lattice vertex algebras.** Suppose σ is an automorphism of $\mathfrak{h}$ with finite order r which preserves the bilinear form $(\cdot|\cdot)$. Then σ is naturally extended to both $\hat{\mathfrak{h}}$ and $M(1)$, again denoted σ. The *σ-twisted Heisenberg algebra* $\hat{\mathfrak{h}}_\sigma$ is spanned over $\mathbb{C}$ by K and the elements $a_m = at^m$ such that $\sigma a = e^{-2\pi \mathrm{i} m} a$. It becomes a Lie algebra with bracket (2.4) for $m, n \in \frac{1}{r}\mathbb{Z}$.

Let $\hat{\mathfrak{h}}_\sigma^{\geq}$ (respectively, $\hat{\mathfrak{h}}_\sigma^{<}$) be the abelian subalgebra of $\hat{\mathfrak{h}}_\sigma$ spanned by all elements a_m with $m \geq 0$ (respectively, $m < 0$). We let $\hat{\mathfrak{h}}_\sigma^{\geq}$ act on $\mathbb{C}$ trivially and K act as the identity operator. The *σ-twisted Fock space* is defined as the induced module

$$M(1)_\sigma = \mathrm{Ind}_{\hat{\mathfrak{h}}_\sigma^{\geq} \oplus \mathbb{C}K}^{\hat{\mathfrak{h}}_\sigma} \mathbb{C} \cong S(\hat{\mathfrak{h}}_\sigma^{<}) \,, \tag{2.10}$$

and is an irreducible highest-weight representation of $\hat{\mathfrak{h}}_\sigma$ which has the structure of a σ-twisted representation of the vertex algebra $M(1)$ (see [FLM, KRR]). For the map Y, we let $Y(\mathbf{1}, z)$ be the identity operator,

$$Y(a,z) = \sum_{n \in p+\mathbb{Z}} a_n \, z^{-n-1} \,, \qquad a \in \mathfrak{h} \,, \;\; \sigma a = e^{-2\pi \mathrm{i} p} a \,, \;\; p \in \frac{1}{r}\mathbb{Z} \,,$$

and extend linearly to all of $\mathfrak{h}$.

Now consider σ as an automorphism of Q. Since σ preserves the bilinear form, the uniqueness of the cocycle ε and (2.7) imply that

$$\eta(\alpha+\beta)\varepsilon(\sigma\alpha, \sigma\beta) = \eta(\alpha)\eta(\beta)\varepsilon(\alpha,\beta) \tag{2.11}$$

for some function $\eta \colon Q \to \{\pm 1\}$. It is shown in [BE, Lemma 2.3] that we can set $\eta = 1$ on any sublattice whose elements satisfy $\varepsilon(\sigma\alpha, \sigma\beta) = \varepsilon(\alpha, \beta)$. In particular, η can be chosen such that

$$\eta(\alpha) = 1 \,, \qquad \alpha \in Q \cap \mathfrak{h}^\sigma \,, \tag{2.12}$$

where $\mathfrak{h}^\sigma \subset \mathfrak{h}$ denotes the subspace spanned by vectors fixed under σ. Then σ can be lifted to an automorphism of the lattice vertex algebra V_Q by setting

$$\sigma(a_n) = \sigma(a)_n \,, \quad \sigma(e^\alpha) = \eta(\alpha) e^{\sigma\alpha} \,, \qquad a \in \mathfrak{h} \,, \; \alpha \in Q \,. \tag{2.13}$$

Note the order of σ on V_Q may double.

It is shown in [BK] that irreducible σ-twisted V_Q-modules are parameterized by the set $(Q^*/Q)^\sigma$ of σ-invariants in Q^*/Q and every σ-twisted V_Q-module is a direct sum of irreducible ones (see [BK, Theorem 4.2]). In the special case when $\sigma = 1$, we get Dong's Theorem that the irreducible V_Q-modules are classified by Q^*/Q (see [D1]). Explicitly, they are given by:

$$V_{\lambda+Q} = \mathcal{F} \otimes \mathbb{C}_\varepsilon[Q] e^\lambda \,, \qquad \lambda \in Q^* \,.$$

When the lattice Q is written as an orthogonal direct sum of sublattices, $Q = L_1 \oplus L_2$, we have a natural isomorphism $V_Q \cong V_{L_1} \otimes V_{L_2}$. It can be shown that if L_1 and L_2 are σ-invariant, there is a correspondence of irreducible twisted modules in the sense that every irreducible σ-twisted V_Q-module M is a tensor product, $M \cong M_1 \otimes M_2$, where M_i is an irreducible $\sigma|_{L_i}$-twisted V_{L_i}-module (cf. [FHL, BE]).

2.4. **The case $\sigma = -1$.** Now we review what is known in the case when $\sigma = -1$, which will be used in our treatment of the general case. In this subsection, we denote the even integral lattice by L instead of Q so that $\mathfrak{h} = \mathbb{C} \otimes_{\mathbb{Z}} L$ is the corresponding complex vector space.

Let $\hat{L}$ be the central extension of L by the 2-group $\langle -1 \rangle$:

$$1 \to \langle -1 \rangle \to \hat{L} \xrightarrow{-} L \to 1 \tag{2.14}$$

with the group structure in $\hat{L}$ written multiplicatively. Then ϵ is the associated 2-cocycle (cf. (2.6)). Let

$$K = \{\sigma(a)a^{-1} \,|\, a \in \hat{L}\} \tag{2.15}$$

and T be any $\hat{L}/K$-module with the natural action of -1. Then these irreducible $\hat{L}/K$-modules are determined by central characters χ of $\hat{L}/K$ such that $\chi(-1) = -1$. When convenient, we denote the corresponding module by T_χ. Define the vector space $V_L^T = M(1)_\sigma \otimes T$ (cf. (2.10)). Then every irreducible σ-twisted V_L-module is isomorphic to V_L^T for some irreducible $\hat{L}/K$-module T_χ (see [FLM, D2]).

An action of σ on V_L^T can be defined by

$$\sigma\left(h^1_{(-n_1)} \cdots h^k_{(-n_k)} t\right) = (-1)^k h^1_{(-n_1)} \cdots h^k_{(-n_k)} t \tag{2.16}$$

for $h^i \in \mathfrak{h}$, $n_i \in \frac{1}{2} + \mathbb{Z}_{\geq 0}$ and $t \in T$. Then σ is an automorphism of V_L^σ-modules. The eigenspaces for σ are denoted $V_L^{T,\pm}$ and are both V_L^σ-modules. Similarly, an action of σ on the untwisted V_L-modules $V_{\lambda+L}$ can be defined by

$$\sigma\left(h^1_{(-n_1)} \cdots h^k_{(-n_k)} e^{\lambda+\alpha}\right) = (-1)^k h^1_{(-n_1)} \cdots h^k_{(-n_k)} e^{-\lambda-\alpha} \tag{2.17}$$

for $h^i \in \mathfrak{h}$, $n_i \in \mathbb{Z}_{\geq 0}$ and $\alpha \in L$. Clearly $\sigma V_{\lambda+L} \subseteq V_{-\lambda+L}$ which implies the eigenspaces $V^\pm_{\lambda+L}$ are V_L^σ-modules for $\lambda \in L^*$ with $2\lambda \in L$ and $V_{\lambda+L} \cong V_{-\lambda+L}$ as V_L^σ-modules if $2\lambda \notin L$. The following theorem is the classification of irreducible V_L^σ-modules.

Theorem 2.1 ([DN, AD])**.** *Let L be a positive-definite even lattice and $\sigma = -1$ on L. Then any irreducible admissible V_L^σ-module is isomorphic to one of the following*:

$$V^\pm_{\lambda+L} \quad (\lambda \in L^*,\, 2\lambda \in L), \quad V_{\lambda+L} \quad (\lambda \in L^*,\, 2\lambda \notin L), \quad V_L^{T,\pm},$$

where T is an irreducible $\hat{L}/K$-module.

Now we describe intertwining operators between the irreducible V_L^σ-modules. For a vector space U, denote by

$$U\{z\} = \Big\{ \sum_{n\in\mathbb{Q}} v_{(n)} z^{-n-1} \Big| v_{(n)} \in U \Big\}$$

the space of U-valued formal series involving rational powers of z. Let V be a vertex algebra, and M_1, M_2, M_3 be V-modules, which are not necessarily distinct. Recall from [FHL] that an *intertwining operator* of type $\binom{M_3}{M_1\ M_2}$ is a linear map given by

$$\mathcal{Y}\colon M_1 \to \mathrm{Hom}(M_2, M_3)\{z\}\,,$$
$$v \mapsto \mathcal{Y}(v,z) = \sum_{n\in\mathbb{Q}} v_{(n)} z^{-n-1}\,, \quad v_{(n)} \in \mathrm{Hom}(M_2, M_3)$$

which satisfies $v_{(n)}u = 0$, for $n \gg 0$, and the Borcherds identity (2.3) for $u \in V, v \in M_1$, where the action of each term is on M_2. The intertwining operators of type $\binom{M_k}{M_i\ M_j}$ form a vector space denoted $\mathcal{V}_{ij}^k$ and the *fusion rule* associated with an algebra V and its modules M_i, M_j, M_k is $N_{i,j}^k = \dim \mathcal{V}_{ij}^k$.

An *admissible* σ-twisted V-module M is a $\frac{1}{r}\mathbb{Z}$-graded V-module

$$M = \bigoplus_{n\in\frac{1}{r}\mathbb{Z}} M(n) \tag{2.18}$$

for which $(v_m M)(n) \subset M(\mathrm{wt}v - m - 1 + n)$ for homogeneous $v \in V$ and $m, n \in \frac{1}{r}\mathbb{Z}$. The *contragredient* M' is defined as the graded dual space

$$M' = \bigoplus_{n\in\frac{1}{r}\mathbb{Z}} M(n)^*, \tag{2.19}$$

where $M(n)^* = \mathrm{Hom}_{\mathbb{C}}(M(n), \mathbb{C})$. A V-module M is *self-dual* if $M = M'$. It is shown in [FHL] that M' has a vertex algebra module structure and the module M is irreducible if and only if M' is irreducible. It is also shown in [FHL] that the fusion rules satisfy a few symmetry properties:

Proposition 2.2 ([FHL])**.** *For V-modules W_1, W_2 and W_3, we have*

$$N_{1,2}^3 = N_{2,1}^3, \quad N_{1,2}^3 = N_{1,3'}^{2'}, \tag{2.20}$$

where prime denotes contragredient module.

The fusion rules for V_L^σ and its irreducible modules were calculated in [A1, ADL] to be either zero or one. In order to present their theorem, we first introduce some additional notation. For $\lambda \in L^*$ such that $2\lambda \in L$, let

$$\pi_{\lambda,\mu} = (-1)^{|\lambda|^2|\mu|^2}, \qquad \lambda, \mu \in L^*, \tag{2.21}$$

$$c_\chi(\lambda) = (-1)^{(\lambda|2\lambda)}\varepsilon(\lambda, 2\lambda)\chi(e^{2\lambda}). \tag{2.22}$$

For any central character χ of $\hat{L}/K$, also set $\chi^{(\lambda)}$ and χ' to be the central characters defined by

$$\chi^{(\lambda)}(a) = (-1)^{(\lambda|\bar{a})}\chi(a), \tag{2.23}$$

$$\chi'(a) = (-1)^{\frac{1}{2}(\bar{a}|\bar{a})}\chi(a), \tag{2.24}$$

for a in the center of $\hat{L}/K$, and set $T_\chi^{(\lambda)} = T_{\chi^{(\lambda)}}$. Note that $\chi' = \chi$ whenever 4 divides $(\bar{a}|\bar{a})$ for all $a \in \hat{L}$. It is shown in [ADL] that

$$(V_{\lambda+L})' \cong V_{\lambda+L}, \quad 2\lambda \notin L \tag{2.25}$$

$$(V_{\lambda+L}^\pm)' \cong \begin{cases} V_{\lambda+L}^\pm, & 2|\lambda|^2 \in 2\mathbb{Z} \\ V_{\lambda+L}^\mp, & 2|\lambda|^2 \in 2\mathbb{Z}+1 \end{cases}, \quad 2\lambda \in L \tag{2.26}$$

$$(V_L^{T_\chi,\pm})' \cong V_L^{T_{\chi'},\pm}. \tag{2.27}$$

The following theorem is only a part of Theorem 5.1 from [ADL].

Theorem 2.3 ([ADL])**.** *Let L be a positive-definite even lattice, and $\epsilon \in \{\pm\}$. Then for two irreducible V_L^σ-modules M_2, M_3, we have the following fusion rules:*

(1) If $\lambda \in L^* \cap \frac{1}{2}L$, the fusion rule of type $\binom{M_3}{V_{\lambda+L}^\epsilon \; M_2}$ is equal to 1 if and only if the pair (M_2, M_3) is one of the following:

$(V_{\mu+L}, V_{\lambda+\mu+L})$, $\mu \in L^*$, $2\mu \notin L$,

$(V_{\mu+L}^{\epsilon_1}, V_{\lambda+\mu+L}^{\epsilon_2})$, $\mu \in L^*$, $2\mu \in L$, $\epsilon_1 \in \{\pm\}$, $\epsilon_2 = \epsilon_1\epsilon\pi_{\lambda,2\mu}$,

$(V_L^{T_\chi,\epsilon_1}, V_L^{T_\chi^{(\lambda)},\epsilon_2})$, $\epsilon_1 \in \{\pm\}$, $\epsilon_2 = c_\chi(\lambda)\epsilon_1\epsilon$.

(2) The fusion rule of type $\binom{M_3}{V_L^{T_\chi,\epsilon} \; M_2}$ is equal to 1 if and only if the pair (M_2, M_3) is one of the following:

$((V_L^{T_\chi^{(\lambda)},\pm})', V_{\lambda+L})$, $(V_{\lambda+L}, V_L^{T_\chi^{(\lambda)},\pm})$, $\lambda \notin L^* \cap \frac{1}{2}L$,

$((V_L^{T_\chi^{(\lambda)},\epsilon_1})', (V_{\lambda+L}^{\epsilon_2})')$, $(V_{\lambda+L}^{\epsilon_2}, V_L^{T_\chi^{(\lambda)},\epsilon_1})$, $\lambda \in L^* \cap \frac{1}{2}L$,

where $\epsilon_1 \in \{\pm\}$, *and* $\epsilon_2 = c_\chi(\lambda)\epsilon_1\epsilon$. *In all other cases, the fusion rules of types* $\binom{M_3}{V^{\epsilon}_{\lambda+L}\ M_2}$ *and* $\binom{M_3}{V_L^{T_\chi,\epsilon}\ M_2}$ *are zero.*

2.5. **Quantum dimension and fusion product.** The notions and properties of quantum dimensions have been systematically studied in [DJX]. A vertex algebra is *rational* if the admissible module category is semisimple (cf. (2.18)). Let $C_2(V) \subset V$ denote the subalgebra spanned by all -2-products. A vertex algebra is C_2-*cofinite* if $V/C_2(V)$ is finite dimensional. Suppose $V = \bigoplus_{n\in\mathbb{Z}} V_n$ is a self dual vertex operator algebra of *CFT type*, i.e., $V_n = 0$ for $n < 0$, and $V_0 = \mathbb{C}\mathbf{1}$. In the case when V is also rational and C_2-cofinite, quantum dimensions of its irreducible modules have nice properties and turn out to be helpful in determining their fusion products.

For a vertex algebra V and a $\frac{1}{r}\mathbb{Z}$-graded σ-twisted V-module M, the *formal character* of M is defined as

$$\mathrm{ch}_q M = q^{\lambda - c/24} \sum_{n\in\frac{1}{r}\mathbb{Z}} (\dim M_{\lambda+n}) q^n, \tag{2.28}$$

where λ is the conformal weight of M. The *quantum dimension* of M over V is defined as

$$q\dim_V M = \lim_{q\to 1^-} \frac{\mathrm{ch}_q M}{\mathrm{ch}_q V}. \tag{2.29}$$

Next we describe the fusion product of two irreducible V-modules, W_1 and W_2. Let $I \in \mathcal{V}^W_{W_1\,W_2}$ be an intertwining operator. A module (W, I) is a *fusion product* of W_1 and W_2 if for any V-module M and $\mathcal{Y} \in \mathcal{V}^M_{W_1\,W_2}$, there is a unique V-module homomorphism $\phi : W \longrightarrow M$ such that $\mathcal{Y} = \phi \circ I$. The fusion product (W, I) is typically denoted by $W_1 \boxtimes W_2$ and in the case when V is rational, it is known that the fusion product exists:

$$W_1 \boxtimes W_2 = \sum_W N^W_{W_1,\,W_2}\, W, \tag{2.30}$$

where the sum runs over all equivalence classes of irreducible V-modules. If V is a simple vertex algebra, a simple V-module M is a *simple current* if for any irreducible V-module W, the fusion product $W \boxtimes M$ exists and is also a simple V-module.

Now assume V is a rational, C_2-cofinite, self dual vertex algebra of CFT type with finitely many irreducible admissible modules. Following [DJX], set $M_0, \ldots, M_d$ as the inequivalent irreducible V-modules with $M_0 \cong V$. Under these conditions, the following properties of quantum dimension are shown in [DJX].

Proposition 2.4 ([DJX]). *(1) $q\dim_V M_i \geq 1$ for any $0 \leq i \leq d$.*
(2) If $q\dim_V M$ exists, then $q\dim_V M = q\dim_V M'$.
(3) For any $0 \leq i, j \leq d$,

$$q\dim_V (M_i \boxtimes M_j) = q\dim_V M_i \cdot q\dim_V M_j.$$

(4) M is a simple current of V if and only if $q\dim_V M = 1$.

We also list a few other basic properties of quantum dimension needed in the calculations.

Lemma 2.5. *Let A, A_i, and B be V-modules for which $A \subset B$ and i is in some index set. Then*

$$q\dim_V A \leq q\dim_V B, \tag{2.31}$$

$$q\dim_V \bigoplus_i A_i = \sum_i q\dim_V A_i. \tag{2.32}$$

Lemma 2.6. *For vertex algebras V and W, let M be a V-module and N be a W-module. Then*

$$q\dim_{V\otimes W} M \otimes N = q\dim_V M \cdot q\dim_W N. \tag{2.33}$$

3. Irreducible V_Q^σ-modules

From now on, we assume σ has order 2. The map σ is extended by linearity to the complex vector space $\mathfrak{h} = \mathbb{C} \otimes_{\mathbb{Z}} Q$ and we denote by

$$\pi_\pm = \frac{1}{2}(1 \pm \sigma), \qquad \alpha_\pm = \pi_\pm(\alpha) \tag{3.1}$$

the projections onto the eigenspaces of σ. Introduce the important sublattices

$$L_\pm = \mathfrak{h}_\pm \cap Q, \qquad L = L_+ \oplus L_- \subseteq Q, \tag{3.2}$$

where $\mathfrak{h}_\pm = \pi_\pm(\mathfrak{h})$. Note that $\mathfrak{h} = \mathfrak{h}_+ \oplus \mathfrak{h}_-$ is an orthogonal direct sum.

In [BE] we constructed and classified the irreducible V_Q^σ-modules and as a consequence of our result [BE, Theorem 3.7], we realized all of them as submodules of twisted or untwisted V_Q-modules [BE, Theorem 3.8]. For a positive-definite even lattice Q and an order two automorphism σ of Q, the first step in our construction is to restrict Q to the sublattice $\bar{Q}$ satisfying $(\alpha|\sigma\alpha) \in 2\mathbb{Z}$ for all $\alpha \in Q$. Under this condition, $\sigma^2 = 1$ on the lattice vertex algebra $V_{\bar{Q}}$, and $(V_Q)^{\sigma^2} = V_{\bar{Q}}$. Therefore

$$V_Q^\sigma = \left((V_Q)^{\sigma^2}\right)^\sigma = V_{\bar{Q}}^\sigma \tag{3.3}$$

so that we may assume $|\sigma| = 2$ on V_Q and only work with the sublattice $\bar{Q}$. For simplicity, we use Q instead of $\bar{Q}$ for the rest of this work. It is

also shown in [BE] that V_Q^σ decomposes as a direct sum of irreducible V_L^σ-modules:

$$V_Q^\sigma \cong \bigoplus_{\gamma+L\in Q/L} V_{\gamma_++L_+}\otimes V^{\eta(\gamma)}_{\gamma_-+L_-}\,. \tag{3.4}$$

Theorem 3.1 ([BE])**.** *Let Q be a positive-definite even lattice, and σ be an automorphism of Q of order two such that $(\alpha|\sigma\alpha)$ is even for all $\alpha\in Q$. Then as a module over $V_L^\sigma\cong V_{L_+}\otimes V^+_{L_-}$ each irreducible V_Q^σ-module is isomorphic to one of the following*:

$$\bigoplus_{\gamma+L\in Q/L} V_{\gamma_++\lambda+L_+}\otimes V_{\gamma_-+\mu+L_-} \qquad (2\mu\notin L_-)\,, \tag{3.5}$$

$$\bigoplus_{\gamma+L\in Q/L} V_{\gamma_++\lambda+L_+}\otimes V^{\epsilon\eta(\gamma)}_{\gamma_-+\mu+L_-} \qquad (2\mu\in L_-)\,, \tag{3.6}$$

where $\lambda\in L_+^*$, $\mu\in L_-^*$, $\epsilon\in\{\pm\}$,

$$\bigoplus_{\gamma+L\in Q/L} V_{\gamma_++\lambda+L_+}\otimes V_{L_-}^{T_\chi^{(\gamma_-)},\epsilon_\gamma} \qquad \epsilon_\gamma=\epsilon\eta(\gamma)c_\chi(\gamma_-)\,, \tag{3.7}$$

where $\lambda\in(\pi_+Q)^*$, and χ is a central character of the group $\hat{L_-}/K$ associated to L_- (c.f. (2.14), (2.15)). Furthermore, every irreducible V_Q^σ-module is a submodule of a V_Q-module or a σ-twisted V_Q-module.

Remark 3.2. The proof is given in [BE]. The only difference is the condition on λ in (3.7). Note that $(\pi_+Q)^*/\pi_+Q\subseteq L_+^*/L_+$ and set

$$v_\gamma=e^\gamma+\eta(\gamma)e^{\sigma\gamma}=e^{\gamma_+}\otimes(e^{\gamma_-}+\eta(\gamma)e^{-\gamma_-})\in V_{\gamma_++L_+}\otimes V^{\eta(\gamma)}_{\gamma_-+L_-}\,. \tag{3.8}$$

Then by the fusion rules in [ADL] we have

$$Y(v_\gamma,z):e^\lambda\otimes V_{L_-}^{T_\chi,\varepsilon}\longrightarrow Y(e^{\gamma_+},z)e^\lambda\otimes V_{L_-}^{T_\chi^{(\gamma_-)},\varepsilon_\gamma}, \tag{3.9}$$

where $Y(e^{\gamma_+},z)e^\lambda = z^{(\gamma_+|\lambda)}\exp\Big(\sum_{n<0}\alpha_n\frac{z^{-n}}{-n}\Big)e^{\gamma_++\lambda}$. To be an untwisted module implies that $(\gamma_+|\lambda)\in\mathbb{Z}$ for all $\gamma\in Q$, i.e., $\lambda\in(\pi_+Q)^*$. Using a similar argument, we also have the restriction that $\lambda+\mu\in Q^*$ in (3.5) and (3.6).

In order to present the fusion products of irreducible V_Q^σ-modules, it is necessary to have more convenient notation. We will denote the modules (3.5) by $U_{\lambda,\mu}$ and refer to them as modules of type 1. Similarly the modules (3.6) are denoted $U^\pm_{\lambda,\mu}$ and we refer to them as modules of type 2. Lastly, the modules (3.7) are denoted $T^{\epsilon_0}_{\lambda,\chi}$, where the sign ϵ_0 is the eigenvalue of σ on the term with $\gamma=0$, and we refer to them as modules of twisted type.

4. Quantum Dimension of Irreducible V_Q^σ-modules

The first important step is to reduce the quantum dimension over V_Q^σ to the quantum dimension over the subalgebra V_L^+.

Proposition 4.1. *We have*

$$q\dim_{V_L^+} V_Q^\sigma = |Q/L|\,, \tag{4.1}$$

and the quantum dimension of any irreducible V_Q^σ-module W can be determined from the quantum dimension of W as a V_L^+-module. In particular,

$$q\dim_{V_Q^\sigma} W = |Q/L|^{-1}\, q\dim_{V_L^+} W. \tag{4.2}$$

Proof. From the fusion rules of irreducible modules of V_{L_+} and $V_{L_-}^+$, we have that $V_{\gamma_+ + L_+} \otimes V^{\eta(\gamma)}_{\gamma_- + L_-}$ is a simple current for every $\gamma \in Q$. The result (4.1) then follows since V_Q^σ has precisely $|Q/L|$ summands (cf. (3.4)). Now let W be an irreducible V_Q^σ-module. Then

$$q\dim_{V_Q^\sigma} W = \frac{q\dim_{V_L^+} W}{q\dim_{V_L^+} V_Q^\sigma} = |Q/L|^{-1}\, q\dim_{V_L^+} W.$$

□

The quantum dimensions for untwisted type irreducible V_Q^σ-modules can now be determined using (4.2) and turn out to be independent of the lattice Q.

Theorem 4.2. *Let W be an irreducible V_Q^σ-module of untwisted type. Then we have the following quantum dimensions:*

$$q\dim_{V_Q^\sigma} W = 2, \quad \text{if } W \text{ is of type 1}, \tag{4.3}$$

$$q\dim_{V_Q^\sigma} W = 1, \quad \text{if } W \text{ is of type 2}. \tag{4.4}$$

In particular, the irreducible V_Q^σ-modules of type 2 are simple currents.

Proof. Let $W = \bigoplus_{\gamma+L\in Q/L} V_{\gamma_+ + \lambda + L_+} \otimes V_{\gamma_- + \mu + L_-}$ be a V_Q^σ-module of type 1 (cf. (3.5)). Using (2.32) and (2.33), we have

$$\begin{aligned} q\dim_{V_L^+} W &= q\dim_{V_L^+} \bigoplus_{\gamma+L\in Q/L} V_{\gamma_+ + \lambda + L_+} \otimes V_{\gamma_- + \mu + L_-} \\ &= \sum_{\gamma+L\in Q/L} q\dim_{V_{L_+}\otimes V_{L_-}^+} V_{\gamma_+ + \lambda + L_+} \otimes V_{\gamma_- + \mu + L_-} \\ &= \sum_{\gamma+L\in Q/L} q\dim_{V_{L_+}} V_{\gamma_+ + \lambda + L_+} \cdot q\dim_{V_{L_-}^+} V_{\gamma_- + \mu + L_-}. \end{aligned}$$

By the fusion rules of irreducible modules of V_{L_+}, the simple module $V_{\gamma_+ + \lambda + L_+}$ is a simple current for V_{L_+} so that

$$q\dim_{V_{L_+}} V_{\gamma_+ + \lambda + L_+} = 1. \tag{4.5}$$

Similarly, $q\dim_{V_{L_-}} V_{\gamma_- + \mu + L_-} = 1$. By the fusion rules of irreducible modules of $V_{L_-}^+$, both simple modules $V_{L_-}^\pm$ are simple currents for $V_{L_-}^+$ so that

$$q\dim_{V_{L_-}^+} V_{L_-} = q\dim_{V_{L_-}^+} V_{L_-}^+ + q\dim_{V_{L_-}^+} V_{L_-}^- = 2. \tag{4.6}$$

Since V_{L_-} is a $V_{L_-}^+$-module, we can write

$$q\dim_{V_{L_-}} V_{\gamma_- + \mu + L_-} = \frac{q\dim_{V_{L_-}^+} V_{\gamma_- + \mu + L_-}}{q\dim_{V_{L_-}^+} V_{L_-}}, \tag{4.7}$$

and therefore

$$q\dim_{V_{L_-}^+} V_{\gamma_- + \mu + L_-} = q\dim_{V_{L_-}} V_{\gamma_- + \mu + L_-} \cdot q\dim_{V_{L_-}^+} V_{L_-} = 2. \tag{4.8}$$

Hence $q\dim_{V_L^+} W = 2|Q/L|$ and $q\dim_{V_Q^\sigma} W = 2$ using (4.2).

Now let $W = \bigoplus_{\gamma + L \in Q/L} V_{\gamma_+ + \lambda + L_+} \otimes V_{\gamma_- + \mu + L_-}^{\epsilon_\gamma}$ be a V_Q^σ-module of type 2, where each $\epsilon_\gamma \in \{\pm\}$ (cf. (3.6)). By fusion rules of irreducible modules of $V_{L_-}^+$, both simple modules $V_{\gamma_- + \mu + L_-}^\pm$ are simple currents. Then similarly by (2.32), (2.33),

$$\begin{aligned} q\dim_{V_L^+} W &= \sum_{\gamma + L \in Q/L} q\dim_{V_{L_+}} V_{\gamma_+ + \lambda + L_+} \cdot q\dim_{V_{L_-}^+} V_{\gamma_- + \mu + L_-}^{\epsilon_\gamma} \\ &= \sum_{\gamma + L \in Q/L} 1 = |Q/L|. \end{aligned}$$

Hence $q\dim_{V_Q^\sigma} W = 1$ by (4.2). □

Quantum dimensions for irreducible V_Q^σ-modules of twisted type can also be realized in terms of quantum dimension of twisted type $V_{L_-}^+$-modules (cf. Proposition 4.1). In order to write this relationship explicitly, we require a closer look at the fusion rules of twisted type modules in [ADL], from which we get the following fusion product:

$$V_{L_-}^{T_\chi,\pm} \boxtimes (V_{L_-}^{T_\chi,\pm})' = \sum_{\substack{2\mu \notin L_- \\ \chi^{(\mu)} = \chi}} V_{\mu + L_-} + \sum_{\substack{2\mu \in L_- \\ \chi^{(\mu)} = \chi}} V_{\mu + L_-}^{\epsilon_\mu}, \tag{4.9}$$

where each $\epsilon_\mu \in \{\pm\}$. We first show that the quantum dimension corresponding to χ and $\chi^{(\gamma_-)}$ are the same for all $\gamma \in Q/L$. In order to use (4.9), we must investigate the solutions to the character equation $\chi^{(\mu)} = \chi$. The condition for the summations in (4.9) for $\chi^{(\gamma_-)}$

is $\left(\chi^{(\gamma_-)}\right)^{(\mu)} = \chi^{(\gamma_-)}$, and this is equivalent to the character equation $\chi^{(\gamma_-+\mu)} = \chi^{(\gamma_-)}$:

$$\begin{aligned}\left(\chi^{(\gamma_-)}\right)^{(\mu)}(U_\alpha) &= (-1)^{(\mu|\alpha)}\chi^{(\gamma_-)}(U_\alpha)\\ &= (-1)^{(\mu+\gamma_-|\alpha)}\chi(U_\alpha)\\ &= \chi^{(\gamma_-+\mu)}(U_\alpha).\end{aligned}$$

It follows that there is a one-to-one correspondence between solutions to the character equations $\chi^{(\gamma_-+\mu)} = \chi^{(\gamma_-)}$ and $\chi^{(\mu)} = \chi$. Using this correspondence, we then obtain:

Lemma 4.3. *The quantum dimensions for the irreducible twisted type $V_{L_-}^+$-modules $V_{L_-}^{T_\chi^{(\gamma_-)},\pm}$ are constant over $\gamma \in Q/L$.*

Proposition 4.4. *Let W be an irreducible V_Q^σ-module of twisted type. Then*

$$q\dim_{V_Q^\sigma} W = q\dim_{V_{L_-}^+} V_{L_-}^{T_\chi,\pm}. \tag{4.10}$$

Proof. Let $W = \bigoplus_{\gamma+L\in Q/L} V_{\gamma_++\lambda+L_+} \otimes V_{L_-}^{T_\chi^{(\gamma_-)},\epsilon_\gamma}$ be a V_Q^σ-module of twisted type, where each $\epsilon_\gamma \in \{\pm\}$ (cf. (3.7)). Then using (2.32), (2.33), (4.5), (4.2), and Lemma 4.3 we have

$$\begin{aligned}q\dim_{V_Q^\sigma} W &= |Q/L|^{-1}\, q\dim_{V_L^+} W\\ &= |Q/L|^{-1}\sum_{\gamma+L\in Q/L} q\dim_{V_{L_+}} V_{\gamma_++\lambda+L_+}\cdot q\dim_{V_{L_-}^+} V_{L_-}^{T_\chi^{(\gamma_-)},\epsilon_\gamma}\\ &= |Q/L|^{-1}\sum_{\gamma+L\in Q/L} q\dim_{V_{L_-}^+} V_{L_-}^{T_\chi^{(\gamma_-)},\epsilon_\gamma}\\ &= q\dim_{V_{L_-}^+} V_{L_-}^{T_\chi,\epsilon_0}.\end{aligned}$$

□

Since $q\dim_{V_{L_-}^+} V_{L_-}^{T_\chi,\pm} = q\dim_{V_{L_-}^+}(V_{L_-}^{T_\chi,\pm})'$, the quantum dimensions of twisted type irreducible V_Q^σ-modules can be calculated using (4.9), Proposition 4.4, and Theorem 4.2. In view of this, we set

$$M = L_-^* \cap \frac{1}{2}L_-, \tag{4.11}$$

so that the number of type 2 irreducible $V_{L_-}^\sigma$-modules is $|M/L_-|$.

It is clear for any central character χ of $\hat{L_-}/K$ that $\chi^{(\mu)} = \chi$ if and only if $\mu \in 2L_-^*$. It follows from (4.9) and Proposition 4.4 that

$q\dim_{V_Q^\sigma} W$ depends on those elements in $2L_-^*$ which correspond to distinct irreducible $V_{L_-}^\sigma$-modules of untwisted type. We denote this finite subset of $2L_-^*$ by R_σ. Then the number of summands in (4.9) is $|R_\sigma|$ and the number of elements in R_σ corresponding to type 2 irreducible $V_{L_-}^\sigma$-modules which appear in (4.9) is $|R_\sigma \cap M|$. We can now write $q\dim_{V_Q^\sigma} W$ in terms of R_σ and M:

Theorem 4.5. *Let W be an irreducible V_Q^σ-module of twisted type and R_σ, M be as defined above. Then the square of the quantum dimension of W over V_Q^σ is given by*

$$(q\dim_{V_Q^\sigma} W)^2 = 2|R_\sigma| - |R_\sigma \cap M|. \tag{4.12}$$

In particular, $(q\dim_{V_Q^\sigma} W)^2 \in \mathbb{Z}$, and $(q\dim_{V_Q^\sigma} W)^2 = |M|$ if $R_\sigma = M$.

Proof. By Proposition 4.4, it is sufficient to find $q\dim_{V_{L_-}^+} V_{L_-}^{T_\chi,\pm}$. We do this using (4.9) and counting the number of irreducible $V_{L_-}^+$-modules of untwisted type. There are precisely $|R_\sigma| - |R_\sigma \cap M|$ distinct irreducible modules of type 1 in the first sum in (4.9) and $|R_\sigma \cap M|$ distinct irreducible modules of type 2 in the second sum in (4.9). The result now follows from (4.9), Proposition 4.4, and Theorem 4.2. □

Remark 4.6. A special case of (4.12) has been proved recently in [DXY2] where the lattice $Q = L \oplus L$ is the direct sum of two copies of a positive definite even lattice L of general rank and σ is the transposition of the two summands (see also [DXY1] for $\operatorname{rank} L = 1$). In this case they find $(q\dim_{V_Q^\sigma} W)^2 = |L^*/L|$.

5. Fusion Products

Recall the notation $U_{\lambda,\mu}$, $U_{\lambda,\mu}^\pm$, and $T_{\lambda,\chi}^{\epsilon_0}$ in Section 3 for the irreducible V_Q^σ-modules. The following is the other main result about fusion products of irreducible V_Q^σ-modules:

Theorem 5.1. *For any $\lambda \in L_+^*$ and $\mu \in L_-^*$, the following are fusion products for irreducible V_Q^σ-modules given in Theorem 3.1:*

$$U_{\lambda,\mu} \boxtimes T^{\epsilon_0}_{\gamma,\chi} = T^{+}_{\lambda+\gamma,\chi^{(\mu)}} + T^{-}_{\lambda+\gamma,\chi^{(\mu)}} \tag{5.1}$$

$$U^{\epsilon}_{\lambda,\mu} \boxtimes T^{\epsilon_0}_{\gamma,\chi} = T^{\epsilon_\mu}_{\lambda+\gamma,\chi^{(\mu)}}, \quad \epsilon_\mu = \epsilon\epsilon_0 c_\chi(\mu) \tag{5.2}$$

$$U_{\lambda,\mu} \boxtimes U^{\epsilon}_{\lambda',\mu'} = U_{\lambda+\lambda',\mu+\mu'} \tag{5.3}$$

$$U^{\epsilon}_{\lambda,\mu} \boxtimes U^{\epsilon'}_{\lambda',\mu'} = U^{\epsilon\epsilon'}_{\lambda+\lambda',\mu+\mu'} \tag{5.4}$$

$$U_{\lambda,\mu} \boxtimes U_{\lambda',\mu'} = \begin{cases} \sum_\epsilon (U^{\epsilon}_{\lambda+\lambda',\mu+\mu'} + U^{\epsilon}_{\lambda+\lambda',\mu-\mu'}) \\ U_{\lambda+\lambda',\mu+\mu'} + U_{\lambda+\lambda',\mu-\mu'} \\ U_{\lambda+\lambda',\mu+\mu'} + U^{+}_{\lambda+\lambda',\mu-\mu'} + U^{-}_{\lambda+\lambda',\mu-\mu'} \end{cases} \tag{5.5}$$

$$T^{\epsilon_0}_{\lambda,\chi} \boxtimes T^{\epsilon'_0}_{\lambda',\psi} = \sum_{\substack{\chi^{(\mu)}=\psi' \\ 2\mu \notin L_-}} U_{\lambda+\lambda',\mu} + \sum_{\substack{\chi^{(\mu)}=\psi' \\ 2\mu \in L_-}} U^{\epsilon_\mu}_{\lambda+\lambda',\mu} \tag{5.6}$$

where $\epsilon_\mu = \epsilon_0\epsilon'_0 c_\chi(\mu)(-1)^{2|\mu|^2}$ in (5.6), the first case in (5.5) is when $2(\mu \pm \mu') \in L_-$ the second case is when $2(\mu \pm \mu') \notin L_-$, and the third case is when $2(\mu + \mu') \notin L_-$ and $2(\mu - \mu') \in L_-$.

Proof. We first prove (5.1). By the fusion rules for irreducible V_{L_+}-modules in [DL] and irreducible $V_{L_-}^{\pm}$-modules in [ADL], the fusion rules of type $\begin{pmatrix} T^{\pm}_{\lambda+\gamma,\chi^{(\mu)}} \\ U_{\lambda,\mu}\ T^{\epsilon_0}_{\gamma,\chi} \end{pmatrix}$ are nonzero and this yields the sum on the right hand side of (5.1). Using (4.12) and Theorem 4.2 to count quantum dimensions of each side of (5.1), we see that there can be no other nonzero fusion rules used in the fusion product.

Similarly, the fusion rules in [DL, ADL] yield nonzero intertwining operators of types $\begin{pmatrix} T^{\epsilon_\mu}_{\lambda+\gamma,\chi^{(\mu)}} \\ U^{\epsilon}_{\lambda,\mu}\ T^{\epsilon_0}_{\gamma,\chi} \end{pmatrix}$, $\begin{pmatrix} U_{\lambda+\lambda',\mu+\mu'} \\ U^{\epsilon}_{\lambda,\mu}\ U_{\lambda',\mu'} \end{pmatrix}$, $\begin{pmatrix} U^{\epsilon\epsilon'}_{\lambda+\lambda',\mu+\mu'} \\ U^{\epsilon}_{\lambda,\mu}\ U^{\epsilon'}_{\lambda',\mu'} \end{pmatrix}$ and a similar argument as for (5.1) proves the fusion products (5.2)-(5.4). For (5.3), note that $2\mu \notin L_-$ and $2\mu' \in L_-$ implies $U_{\lambda+\lambda,\,\mu+\mu'} \cong U_{\lambda+\lambda,\,\mu-\mu'}$. Also, since $q\dim(U^{\epsilon}_{\lambda,\mu} \boxtimes U^{\epsilon'}_{\lambda',\mu'}) = 1$ by (4.4) and Proposition 2.4, the fusion product $U^{\epsilon}_{\lambda,\mu} \boxtimes U^{\epsilon'}_{\lambda',\mu'}$ is a simple current, and therefore must equal one irreducible module of type 2 (cf. (4.12), Theorem 4.2). We then obtain (5.4), where the given rule of signs follows because $2\mu \in L_-$ implies $\pi_{\lambda,2\mu} = 1$ (cf. (2.21)).

We now prove (5.5) and (5.6). By the fusion rules in [DL, ADL], the intertwining operators of type $\begin{pmatrix} U_{\lambda+\lambda',\mu+\mu'} \\ U_{\lambda,\mu}\ U_{\lambda',\mu'} \end{pmatrix}$ are nonzero only when

$\mu+\mu' \notin 2L_-$ and the intertwining operators of type $\binom{U^{\pm}_{\lambda+\lambda',\,\mu+\mu'}}{U_{\lambda,\mu}\ U_{\lambda',\mu'}}$ are nonzero only when $\mu+\mu' \in 2L_-$. By counting quantum dimensions of both sides of (5.5) we see that there can be no other nonzero fusion rules used in the fusion product for either case. This proves (5.5). Again by fusion rules in [DL, ADL], the intertwining operators of type $\binom{U_{\lambda+\lambda',\mu}}{T^{\epsilon_0}_{\lambda,\chi}\ T^{\epsilon'_0}_{\lambda',\psi}}$ are nonzero only when $\chi^{(\mu)}=\psi'$ and $2\mu \notin L_-$, and the intertwining operators of type $\binom{U^{\epsilon_\mu}_{\lambda+\lambda',\mu}}{T^{\epsilon_0}_{\lambda,\chi}\ T^{\epsilon'_0}_{\lambda',\psi}}$ are nonzero only when $\chi^{(\mu)}=\psi'$ and $2\mu \in L_-$ (cf. (2.24)). In order to count the quantum dimension of the right-hand side of (5.6), we count the number of solutions $\mu+L_-$ to the character equation $\chi^{(\mu)}=\psi'$. Let $R^{\chi,\psi}_\sigma$ be the solution set, where each element is distinct under σ. Clearly then for any $\mu+L_- \in R^{\chi,\psi}_\sigma$, we have $\mu+L_-+R_\sigma \subset R^{\chi,\psi}_\sigma$ (cf. (**??**)). We show these two sets are equal, i.e., that $\mu,\mu' \in R^{\chi,\psi}_\sigma$ implies $\mu-\mu'+L_- \in R_\sigma$:

$$\begin{aligned}
\chi^{(\mu-\mu')}(U_\alpha) &= (-1)^{(\mu-\mu'|\alpha)}\chi(U_\alpha)\\
&= (-1)^{(\mu|\alpha)+(\mu'|\alpha)+(\alpha|\alpha)}\chi(U_\alpha)\\
&= (-1)^{(\mu|\alpha)+\frac{1}{2}(\alpha|\alpha)}(-1)^{(\mu'|\alpha)+\frac{1}{2}(\alpha|\alpha)}\chi(U_\alpha)\\
&= (-1)^{(\mu|\alpha)+\frac{1}{2}(\alpha|\alpha)}\psi(U_\alpha)\\
&= \chi(U_\alpha),
\end{aligned}$$

where the last two steps use $\chi^{(\mu)}=\psi'$. Therefore $|R^{\chi,\psi}_\sigma| = |R_\sigma|$. By counting quantum dimensions of both sides of (5.6) we now see there can be no other nonzero fusion rules used in the fusion product. Finally, the rule for the sign ϵ_μ in (5.6) follows from the fusion rules in Theorem 2.3 and (2.26). This proves (5.6). □

6. Examples

Here we show explicit details of quantum dimensions and fusion products of twisted type orbifold modules in the cases of a 2-permuatation orbifold and for the A_n root lattice with a Dynkin diagram automorphism.

6.1. $(V_K \otimes V_K)^{\mathbb{Z}_2}$ **orbifolds.** The case when Q is an orthogonal direct sum of two copies of an arbitrary rank positive definite even lattice K with the 2-cycle permutation has recently been studied in [DXY2]. Here we present how some of our results can simplify in these cases

and show the explicit results in the special case of the A_2 root lattice. In this section, we use the symbol K to avoid confusion with (3.2).

Let $Q = K \oplus K$ be an orthogonal direct sum, where $K = \langle \alpha_1, \ldots, \alpha_n \rangle$ is a positive definite even lattice with rank n. Set $(\alpha_i|\alpha_i) = 2k_i$ and for $\gamma \in K$, set $\gamma^1 = (\gamma, 0)$ and $\gamma^2 = (0, \gamma)$. Then a 2-cocycle ε on K can be extended to a 2-cocycle on Q for which we may take $\eta = 1$ on Q (cf. (2.13)). Let σ be the 2-cycle permutation $\sigma(a, b) = (b, a)$. Then the vectors

$$\alpha^i = \alpha_i^1 + \alpha_i^2 = (\alpha_i, \alpha_i), \tag{6.1}$$

$$\beta^i = \alpha_i^1 - \alpha_i^2 = (\alpha_i, -\alpha_i), \tag{6.2}$$

for $i = 1, \ldots, n$, are eigenvectors of σ with eigenvalues ± 1 and

$$(\alpha^i|\alpha^i) = 4k_i = (\beta^i|\beta^i), \tag{6.3}$$

$$(\alpha^i|\alpha^j) = 2(\alpha_i|\alpha_j) = (\beta^i|\beta^j). \tag{6.4}$$

These relations then imply $Q = \bar{Q}$ (cf. Section 3).

First we describe the orbifold V_Q^σ. We have (cf. (3.2), (3.4))

$$L_+ = \langle \alpha^1, \ldots, \alpha^n \rangle, \quad L_- = \langle \beta^1, \ldots, \beta^n \rangle, \tag{6.5}$$

$$V_Q^\sigma = \bigoplus_{(b_1,\ldots,b_n)} \left(V_{\frac{1}{2}\sum b_i \alpha^i + L_+} \otimes V^+_{\frac{1}{2}\sum b_i \beta^i + L_-} \right), \tag{6.6}$$

where each $b_i \in \{0, 1\}$ and there are $|Q/L| = 2^n$ summands. We have

$$M = \left\langle \frac{\beta^i}{2} \,\middle|\, i = 0, \ldots, n \right\rangle, \tag{6.7}$$

$$R_\sigma \cap M = \operatorname{span}_{\mathbb{Z}_2} \left\{ \frac{\beta^i}{2} \,\middle|\, (\alpha_i|K) \in 2\mathbb{Z} \right\}, \tag{6.8}$$

and in this case, the Gram matrix for K can determine the size of $R_\sigma \cap M$.

Lemma 6.1. *Let M be as in* (4.11). *Then $|R_\sigma \cap M| = 1$ if and only if for every i, $(\alpha_i|\alpha_j)$ is odd for at least one j.*

Proof. Let $\frac{\beta^i}{2} \in M$ and j be such that $(\alpha_i|\alpha_j)$ is odd. Then

$$\left(\frac{\beta^i}{2} \middle| \beta^j \right) = (\alpha_i|\alpha_j), \qquad \chi^{(\frac{\beta^i}{2})}(e^{\beta^j}) = -\chi(e^{\beta^j}), \tag{6.9}$$

so that $\frac{\beta^i}{2} \notin R_\sigma$. □

Due to (6.3), we have

$$\chi' = \chi \tag{6.10}$$

for all central characters χ of L_-. Hence all irreducible V_Q^σ-modules of twisted type are self dual.

6.1.1. *Rank 1 case.* Here we provide explicit details of Theorem 4.5 in the case when K is a rank 1 even positive definite lattice. These results will correspond to the results previously done in [DXY1].

Write $K = \mathbb{Z}\gamma$, where $(\gamma|\gamma) = 2k$ for some positive integer k and let σ be the 2-cycle permutation. Then it is easy to see that

$$L_- = \mathbb{Z}\beta, \qquad (\beta|\beta) = 4k, \tag{6.11}$$

$$L_-^* = \frac{1}{4k}L_-, \qquad 2L_-^* = \frac{1}{2k}L_-. \tag{6.12}$$

We also have $|R_\sigma \cap M| = 2$ and the other elements of L_-^*/L_- are paired under σ. Therefore the quantum dimension of any twisted type irreducible V_Q^σ-module W is given by Theorem 4.5:

$$(q\dim_{V_Q^\sigma} W)^2 = 2|R_\sigma| - |R_\sigma \cap M| \tag{6.13}$$

$$= 2\left(\frac{2k-2}{2} + 2\right) - 2 \tag{6.14}$$

$$= 2k. \tag{6.15}$$

6.1.2. $A_2 \oplus A_2$ *case.* Here we provide explicit details of Theorems 3.1 and 4.5 in the case when $K = \mathbb{Z}\alpha_1 + \mathbb{Z}\alpha_2$ is the A_2 root lattice.

Consider the orthogonal direct sum $Q = K \oplus K$ and let σ be the 2-cycle permutation. Then

$$L_+ = \langle \alpha^1, \alpha^2 \rangle, \quad L_- = \langle \beta^1, \beta^2 \rangle, \tag{6.16}$$

$$|\alpha^i|^2 = 4 = |\beta^i|^2, \quad (\alpha^1|\alpha^2) = -2 = (\beta^1|\beta^2), \tag{6.17}$$

and the orbifold is given by

$$V_Q^\sigma = \bigoplus_{(b_1,b_2)} \left(V_{\frac{1}{2}\sum b_i\alpha^i + L_+} \otimes V^+_{\frac{1}{2}\sum b_i\beta^i + L_-} \right), \tag{6.18}$$

where $b_1, b_2 \in \{0, 1\}$. To describe the dual lattices $L_\pm^*$ we set

$$\delta^1 = \frac{\alpha^1 + 2\alpha^2}{6}, \quad \delta^2 = \frac{2\alpha^1 + \alpha^2}{6}, \tag{6.19}$$

$$\rho^1 = \frac{\beta^1 + 2\beta^2}{6}, \quad \rho^2 = \frac{2\beta^1 + \beta^2}{6}, \tag{6.20}$$

so that $L_+^* = \langle \delta^1, \delta^2 \rangle$ and $L_-^* = \langle \rho^1, \rho^2 \rangle$ with quotients $L_\pm^*/L_\pm \simeq \mathbb{Z}_2 \times \mathbb{Z}_6$. Note that $\sigma(\rho^i + L_-) = 5\rho^i + L_-, 2\rho^2 + L_- = 4\rho^1 + L_-,$

and $2\delta^2 + L_+ = 4\delta^1 + L_+$. By Theorem 3.1, the type 1 irreducible V_Q^σ-modules are

$$U_{\lambda_{ij},\mu} = \bigoplus_{(b_1,b_2)} \left(V_{\lambda_{ij}+\frac{1}{2}\sum b_i\alpha^i+L_+} \otimes V_{\mu+\frac{1}{2}\sum b_i\beta^i+L_-} \right), \tag{6.21}$$

where $\lambda_{ij} = i\delta^1 + j\delta^2$, $0 \le i \le 5$, $j = 0, 1$ and $\mu \in \{\rho^1, 2\rho^1, \rho^2, \rho^1 + 5\rho^2\}$. Set $\mu = k\rho^1 + l\rho^2$. Then for (6.21) to be untwisted, we also require that $i + k \in 2\mathbb{Z}$ and $j + l \in 2\mathbb{Z}$ (cf. Remark 3.2). This gives 12 distinct irreducible modules of type 1.

Now we find

$$M/L_- = \left\langle 3\rho^1 + L_-, 3\rho^2 + L_- \right\rangle \simeq \mathbb{Z}_2^2, \tag{6.22}$$

(cf. (4.11)) so that the type 2 irreducible V_Q^σ-modules are given by $U^\pm_{\lambda_{ij},\mu}$, with $\mu + L_- \in M/L_-$. In order for $U^\pm_{\lambda_{ij},\mu}$ to be untwisted we also require that $i + k \in 2\mathbb{Z}$ and $j + l \in 2\mathbb{Z}$ (cf. Remark 3.2). This gives 24 distinct irreducible modules of type 2.

For the orbifold modules of twisted type, consider the following labelling of the central characters:

	1	e^{β^1}	e^{β^2}	$e^{\beta^1+\beta^2}$
χ_{00}	1	1	1	1
χ_{10}	1	1	−1	−1
χ_{01}	1	−1	1	−1
χ_{11}	1	−1	−1	1

For $\gamma = c_1\beta^1 + c_2\beta^2$, notice that $\chi_{ij}(e^\gamma) = (-1)^{c_1 j + c_2 i}$. We can then use this labelling to show the following properties:

Lemma 6.2. *Let $b_i, c_i \in \{0, 1\}$ and $\mu = \frac{1}{2}b_1\beta^1 + \frac{1}{2}b_2\beta^2$. Then we have*

$$c_\chi(\mu) = \chi(e^{b_1\beta^1})\chi(e^{b_2\beta^2}), \tag{6.23}$$

$$\chi_{ij}^{(\mu)} = \chi_{i+b_1,j+b_2} \tag{6.24}$$

$$\chi^{(r\rho^1+k\rho^2)}(e^{\vec{c}\cdot\vec{\beta}}) = (-1)^{rc_2+kc_1}\chi(e^{\vec{c}\cdot\vec{\beta}}), \tag{6.25}$$

where $\vec{c}\cdot\vec{\beta} = c_1\beta^1 + c_2\beta^2$ and the indices in (6.24) are taken modulo 2.

Now since $(\pi_+Q)^*/(\pi_+Q) = \langle 2\delta^1 + \pi_+Q\rangle \simeq \mathbb{Z}_3$, there are $3 \cdot 4 = 12$ orbifold modules of twisted type:

$$T^{\epsilon_0}_{2i\delta^1,\chi_{mn}} = \bigoplus_{\mu=(b_1,b_2)} V_{2i\delta^1+\frac{1}{2}\sum b_i\alpha^i+L_+} \otimes V_{L_-}^{T_{\chi_{m+b_1,n+b_2}},\epsilon_\mu}, \tag{6.26}$$

where $i = 0, 1, 2$, $\epsilon_\mu = \epsilon c_\chi(\mu)$ (cf. (6.23)).

Now we determine the quantum dimensions for irreducible modules of twisted type:

$$R_\sigma = \{0, 2\rho^1\}, \quad |R_\sigma \cap M| = 1, \tag{6.27}$$

$$q\dim_{V_Q^\sigma} T^{\epsilon_0}_{\lambda,\chi_{ij}} = \sqrt{3}, \tag{6.28}$$

(cf. Lemma 6.1) and a few of their fusion products:

$$T^{\epsilon_0}_{\lambda,\chi_{00}} \boxtimes T^{\epsilon'_0}_{\lambda',\chi_{10}} = U_{\lambda+\lambda',\rho^1} + U^{\epsilon_0\epsilon'_0}_{\lambda+\lambda',3\rho^1}, \tag{6.29}$$

$$T^{\epsilon_0}_{\lambda,\chi_{00}} \boxtimes T^{\epsilon'_0}_{\lambda',\chi_{01}} = U_{\lambda+\lambda',\rho^2} + U^{\epsilon_0\epsilon'_0}_{\lambda+\lambda',3\rho^2}, \tag{6.30}$$

$$T^{\epsilon_0}_{\lambda,\chi_{00}} \boxtimes T^{\epsilon'_0}_{\lambda',\chi_{11}} = U_{\lambda+\lambda',\rho^1+5\rho^2} + U^{\epsilon_0\epsilon'_0}_{\lambda+\lambda',\rho^1+\rho^2}. \tag{6.31}$$

The other fusion products are similar. Note that each fusion product can be written in the form $U_{\lambda,\mu} + U^{\pm}_{\lambda,\mu'}$ $(\lambda \in L_+,\ \mu, \mu' \in L_-)$, and that $q\dim(T^{\epsilon_0}_{\lambda,\chi_{ij}} \boxtimes T^{\epsilon'_0}_{\lambda',\chi_{kl}}) = 3$ for $i, j, k, l \in \{0, 1\}$ (cf. (6.28)).

6.2. **The A_n root lattice.** Consider the A_n Dynkin diagram with the simple roots $\{\alpha_1, \dots, \alpha_n\}$, the associated root lattice $Q = \sum_{i=1}^n \mathbb{Z}\alpha_i$, and the Dynkin diagram automorphism $\sigma : \alpha_i \leftrightarrow \alpha_{n-i+1}$. Set

$$\alpha^i = \alpha_i + \alpha_{n-i+1} \quad \text{and} \quad \beta^i = \alpha_i - \alpha_{n-i+1}. \tag{6.32}$$

The construction of all irreducible V_Q^σ-modules was done in [E]. Since the results depend on the parity of n, we treat the cases separately.

6.2.1. *Results for n odd.* Throughout this section, set $l = \frac{n-1}{2}$ and let $i < l + 1$. Then

$$|\alpha^i|^2 = 4 = |\beta^i|^2, \quad (\alpha^i|\beta^j) = 0, \quad (\alpha^i|\alpha^{i+1}) = -2 = (\beta^i|\beta^{i+1}),$$

and $(\alpha^i|\alpha^j) = 0 = (\beta^i|\beta^j)$ otherwise, and

$$L_+ \ = \ \sum_{i=1}^{l} \mathbb{Z}\alpha^i + \mathbb{Z}\alpha_{l+1}, \quad L_- = \sum_{i=1}^{l} \mathbb{Z}\beta^i.$$

Therefore $Q = \bar{Q}$ (cf. Section 3), and the orbifold is given by

$$V_Q^\sigma \simeq \bigoplus_{(b_1,\dots,b_l)} \left(V_{\frac{1}{2}\sum b_i\alpha^i + L_+} \otimes V^+_{\frac{1}{2}\sum b_i\beta^i + L_-} \right), \tag{6.33}$$

where $b_1, \dots, b_l \in \{0, 1\}$. Now the Gram matrix for L_- is twice the Gram matrix for A_l so that we may use the dual lattice $\{\lambda_1, \dots, \lambda_l\}$ for A_l to describe the dual lattice for L_-. A basis for L_-^* is then

$$\mu_i = \frac{1}{2(l+1)}((l-i+1)\beta^1 + 2(l-i+1)\beta^2 + i(l-i+1)\beta^i + i(l-i)\beta^{i+1} + \cdots + i\beta^l),$$

where $i = 1, \dots, l$. Then there are $l + 4$ irreducible V_Q^σ-modules of untwisted type (see [E] Theorem 5.5.5):

$$U_0^\pm = \bigoplus_{(b_1,\dots,b_l)} \left(V_{\frac{1}{2}\sum b_i\alpha^i + L_+} \otimes V^\pm_{\frac{1}{2}\sum b_i\beta^i + L_-} \right),$$

$$U_k = \bigoplus_{(b_1,\dots,b_l)} \left(V_{\frac{1}{2}\epsilon_k\alpha_{l+1} + \frac{1}{2}\sum b_i\alpha^i + L_+} \otimes V_{k\mu_l + \frac{1}{2}\sum b_i\beta^i + L_-} \right), \quad k = 1, \dots, l$$

$$U_{l+1}^\pm = \bigoplus_{(b_1,\dots,b_l)} \left(V_{\frac{1}{2}\epsilon_l\alpha_{l+1} + \frac{1}{2}\sum b_i\alpha^i + L_+} \otimes V^\pm_{(l+1)\mu_l + \frac{1}{2}\sum b_i\beta^i + L_-} \right),$$

where $b_1, \dots, b_l \in \{0, 1\}$ and $\epsilon_i = 0$ for i odd and $\epsilon_i = 1$ for i even. From this it is clear that $|R_\sigma| = \left\lfloor \frac{l}{2} \right\rfloor + 2$ and $|R_\sigma \cap M| = 2$. If W is an irreducible V_Q^σ-module of twisted type, then from Theorem 4.5 we have

$$(q \dim_{V_Q^\sigma} W)^2 = \begin{cases} l + 2, & l \text{ even}, \\ l + 1, & l \text{ odd} \end{cases}.$$

6.2.2. *Results for n even.* Throughout this section, set $l = \frac{n}{2}$ and let $i < l + 1$. Then for $i < l$,

$$|\alpha^i|^2 = 4 = |\beta^i|^2, \quad (\alpha^i|\beta^j) = 0, \quad (\alpha^i|\alpha^{i+1}) = -2 = (\beta^i|\beta^{i+1}),$$

$(\beta^l|\beta^l) = 6$, $(\alpha^l|\alpha^l) = 2$, and $(\alpha^i|\alpha^j) = 0 = (\beta^i|\beta^j)$ otherwise. Then

$$\bar{Q} = \sum_{i=1}^{l-1} \mathbb{Z}\alpha_i + \mathbb{Z}\alpha^l + \mathbb{Z}\beta^l + \sum_{i=l+2}^{n} \mathbb{Z}\alpha_i,$$

and the cosets $\bar{Q}/L$ are in correspondence with $\{0, 1\}$-valued $(l - 1)$-tuples (cf. Section 3). Therefore the orbifold is given by

$$V_Q^\sigma \simeq \bigoplus_{(b_1,\dots,b_{l-1})} \left(V_{\frac{1}{2}\sum b_i\alpha^i + L_+} \otimes V^+_{\frac{1}{2}\sum b_i\beta^i + L_-} \right), \tag{6.34}$$

where $b_1, \dots, b_{l-1} \in \{0, 1\}$. A basis of L_-^* was computed in [E, Proposition 5.6.2]:

$$\mu_i = \frac{1}{2(2l+1)}((2l - 2i + 1)\beta^1 + \cdots + i(2l - 2i + 1)\beta^i + i(2l - 2i - 1)\beta^{i+1} + \cdots + i\beta^l),$$

where $1 \le i \le l$. Set $\gamma = \frac{1}{2}(\beta^1 + \beta^3 + \cdots + \beta^s)$, where $s = \begin{cases} l, & l \text{ odd} \\ l - 1, & l \text{ even} \end{cases}$.

Then there are $l+4$ irreducible V_Q^σ-modules of untwisted type (see [E] Theorem 5.6.6):

$$U_0^\pm = \bigoplus_{(b_1,\dots,b_{l-1})} \left(V_{\frac{1}{2}\sum b_i\alpha^i+L_+} \otimes V^\pm_{\frac{1}{2}\sum b_i\beta^i+L_-} \right)$$

$$U_k = \bigoplus_{(b_1,\dots,b_{l-1})} \left(V_{\frac{1}{2}\sum b_i\alpha^i+L_+} \otimes V_{2k\mu_l+\frac{1}{2}\sum b_i\beta^i+L_-} \right), \quad k=1,\dots,l-1,$$

$$U_\gamma^\pm = \bigoplus_{(b_1,\dots,b_{l-1})} \left(V_{\frac{1}{2}\sum b_i\alpha^i+L_+} \otimes V^\pm_{\gamma+\frac{1}{2}\sum b_i\beta^i+L_-} \right),$$

where $b_1,\dots,b_l \in \{0,1\}$. Note that $\frac{\beta^l}{2} \notin R_\sigma$. From this it is clear that

$$|R_\sigma| = \begin{cases} l+1, & l \text{ even}, \\ l, & l \text{ odd} \end{cases}, \quad |R_\sigma \cap M| = \begin{cases} 2, & l \text{ even}, \\ 1, & l \text{ odd} \end{cases}.$$

If W is an irreducible V_Q^σ-module of twisted type, then from Theorem 4.5 we have

$$(q\dim_{V_Q^\sigma} W)^2 = \begin{cases} 2l, & l \text{ even}, \\ 2l-1, & l \text{ odd} \end{cases}.$$

Acknowledgements

The author is grateful to Bojko Bakalov for helpful discussions and revisions of drafts. The author is also grateful to the reviewer for thoughtful comments and suggestions.

References

[A1] T. Abe, *Fusion rules for the charge conjugation orbifold*. J. Algebra **242** (2001), 624–655.

[A2] T. Abe, *Rationality of the vertex operator algebra V_L^+ for a positive definite even lattice L*. Math. Z. **249** (2005), 455–484.

[ABD] T. Abe, G. Buhl, and C. Dong, *Rationality, regularity, and C_2-cofiniteness*. Trans. Amer. Math. Soc. **356** (2004), 3391–3402.

[AD] T. Abe and C. Dong, *Classification of irreducible modules for the vertex operator algebra V_L^+ : general case*. J. Algebra **273** (2004), 657–685.

[ADL] T. Abe, C. Dong, and H. Li, *Fusion rules for the vertex operator algebra $M(1)$ and V_L^+*. Comm. Math. Phys. **253** (2005), 171–219.

[BK] B. Bakalov and V.G. Kac, *Twisted modules over lattice vertex algebras*. In: "Lie theory and its applications in physics V," 3–26, World Sci. Publishing, River Edge, NJ, 2004; math.QA/0402315.

[B] R.E. Borcherds, *Vertex algebras, Kac–Moody algebras, and the Monster*. Proc. Nat. Acad. Sci. USA **83** (1986), 3068–3071.

[BE] B. Bakalov and J. Elsinger, *Orbifolds of lattice vertex algebras under an isometry of order two*. J. Algebra **441** (2015), 57–83.

[DJX] C. Dong, X. Jiao, F. Xu *Quantum dimensions and quantum Galois theory*. Trans. Amer. Math. Soc. **365** (2013), 6441–6469.

[DVVV] R. Dijkgraaf, C. Vafa, E. Verlinde, and H. Verlinde, *The operator algebra of orbifold models*. Comm. Math. Phys. **123** (1989), 485–526.

[D1] C. Dong, *Vertex algebras associated with even lattices*. J. Algebra **161** (1993), 245–265.

[D2] C. Dong, *Twisted modules for vertex algebras associated with even lattices*. J. Algebra **165** (1994), 91–112.

[DJL] C. Dong, C. Jiang, and X. Lin, *Rationality of vertex operator algebra* V_L^+ : *higher rank*. Proc. London Math. Soc. **104** (2012), 799–826.

[DL] C. Dong and J. Lepowsky, *Generalized vertex algebras and relative vertex operators*. Progress in Math., 112, Birkhäuser Boston, 1993.

[DLM1] C. Dong, H. Li, and G. Mason, *Modular-invariance of trace functions in orbifold theory and generalized Moonshine*. Comm. Math. Phys. **214** (2000), 1–56.

[DLM2] C. Dong, H. Li, and G. Mason, *Regularity of rational vertex operator algebra*. Adv. Math. **312** (1997), 148–166.

[DN] C. Dong and K. Nagatomo, *Representations of vertex operator algebra* V_L^+ *for rank* 1 *lattice* L, Comm. Math. Phys. **202** (1999), 169–195.

[DXY1] C. Dong, F. Xu, and N. Yu, *2-permutations of lattice vertex operator algebras: Higher rank*, http://arxiv.org/abs/1602.05602

[DXY2] C. Dong, F. Xu, and N. Yu, *Cyclic permutations of lattice vertex operator algebras*, http://dx.doi.org/10.1090/proc/12966

[E] J.R. Elsinger, *Classification of orbifold modules under an automorphism of order two*. Ph.D. dissertation, 2014, NC State University; http://www.lib.ncsu.edu/resolver/1840.16/9824

[FFR] A.J. Feingold, I.B. Frenkel, and J.F.X. Ries, *Spinor construction of vertex operator algebras, triality, and* $E_8^{(1)}$. Contemporary Math., 121, Amer. Math. Soc., Providence, RI, 1991.

[FB] E. Frenkel and D. Ben-Zvi, *Vertex algebras and algebraic curves*. Math. Surveys and Monographs, 88, Amer. Math. Soc., Providence, RI, 2001; 2nd ed., 2004.

[FHL] I.B. Frenkel, Y.-Z. Huang, and J. Lepowsky, *On axiomatic approaches to vertex operator algebras and modules*. Mem. Amer. Math. Soc. **104** (1993), no. 494.

[FK] I.B. Frenkel and V.G. Kac, *Basic representations of affine Lie algebras and dual resonance models*. Invent. Math. **62** (1980), 23–66.

[FLM] I. B. Frenkel, J. Lepowsky, and A. Meurman, "Vertex operator algebras and the Monster," Pure and Appl. Math., 134, Academic Press, Boston, 1988.

[K] V.G. Kac, *Vertex algebras for beginners*. University Lecture Series, 10, Amer. Math. Soc., Providence, RI, 1996; 2nd ed., 1998.

[KRR] V.G. Kac, A.K. Raina, and N. Rozhkovskaya, *Bombay lectures on highest weight representations of infinite dimensional Lie algebras*. 2nd ed., Advanced Ser. in Math. Phys., 29. World Sci. Pub. Co. Pte. Ltd., Hackensack, NJ, 2013.

[KT] V.G. Kac and I.T. Todorov, *Affine orbifolds and rational conformal field theory extensions of* $W_{1+\infty}$. Comm. Math. Phys. **190** (1997), 57–111.

[LL] J. Lepowsky and H. Li, *Introduction to vertex operator algebras and their representations*. Progress in Math., 227, Birkhäuser Boston, Boston, MA, 2004.

[M1] M. Miyamoto, *A $\mathbb{Z}_3$-orbifold theory of lattice vertex operator algebra and $\mathbb{Z}_3$-orbifold constructions*. In: "Symmetries, integrable systems and representations," 319–344, Springer Proc. Math. Stat., 40, Springer, Heidelberg, 2013; arXiv:1003.0237.

[M2] M. Miyamoto, *C_2-cofiniteness of cyclic-orbifold models*. Preprint arXiv:1306.5031.

[M3] M. Miyamoto, *Flatness and semi-rigidity of vertex operator algebras*. Preprint arXiv:1104.4675.

[MC] M. Miyamoto, S. Carnahan, *Rationality of fixed-point vertex operator algebras*. Preprint arXiv:1603.05645v1.

Department of Mathematics, Spring Hill College, Mobile, AL 36695, USA

E-mail address: jelsinger@shc.edu